



Nicotinamide Deficiency in Primary Open-Angle Glaucoma

Submitted by Stéphanie Pinot on Tue, 06/25/2019 - 15:21

Titre Nicotinamide Deficiency in Primary Open-Angle Glaucoma

Type de publication Article de revue

Auteur Kouassi Nzougnet, Judith [1], Chao de La Barca, Juan Manuel [2], Guehlouz, Khadidja [3], Leruez, Stéphanie [4], Coulbault, Laurent [5], Allouche, Stéphane [6], Bocca, Cinzia [7], Muller, Jeanne [8], Amati-Bonneau, Patrizia [9], Gohier, Philippe [10], Bonneau, Dominique [11], Simard, Gilles [12], Milea, Dan [13], Lenaers, Guy [14], Procaccio, Vincent [15], Reynier, Pascal [16]

Editeur Association for Research in Vision and Ophthalmology

Type Article scientifique dans une revue à comité de lecture

Année 2019

Langue Anglais

Date Juin 2019

Numéro 7

Pagination 2509-2514

Volume 60

Titre de la revue Investigative ophthalmology & visual science

ISSN 1552-5783

Purpose: To investigate the plasma concentration of nicotinamide in primary open-angle glaucoma (POAG).
Methods: Plasma of 34 POAG individuals was compared to that of 30 age- and sex-matched controls using a semiquantitative method based on liquid chromatography coupled to high-resolution mass spectrometry. Subsequently, an independent quantitative method, based on liquid chromatography coupled to mass spectrometry, was used to assess nicotinamide concentration in the plasma from the same initial cohort and from a replicative cohort of 20 POAG individuals and 15 controls.
Results: Using the semiquantitative method, the plasma nicotinamide concentration was significantly lower in the initial cohort of POAG individuals compared to controls and further confirmed in the same cohort, using the targeted quantitative method, with mean concentrations of 0.14 μ M (median: 0.12 μ M; range, 0.06-0.28 μ M) in the POAG group (-30%; $P = 0.022$) and 0.19 μ M (median: 0.18 μ M; range, 0.08-0.47 μ M) in the control group. The quantitative dosage also disclosed a significantly lower plasma nicotinamide concentration (-33%; $P = 0.011$) in the replicative cohort with mean concentrations of 0.14 μ M (median: 0.14 μ M; range, 0.09-0.25 μ M) in the POAG group, and 0.19 μ M (median: 0.21 μ M; range, 0.09-0.26 μ M) in the control group.
Conclusions: Glaucoma is associated with lower plasmatic nicotinamide levels, compared to controls, suggesting that nicotinamide supplementation might become a future therapeutic strategy. Further studies are needed, in larger cohorts, to confirm these preliminary findings.

Résumé en anglais

URL de la notice <http://okina.univ-angers.fr/publications/ua19841> [17]

DOI	10.1167/iovs.19-27099 [18]
Lien vers le document	https://iovs.arvojournals.org/article.aspx?articleid=2736058 [19]
Titre abrégé	Invest. Ophthalmol. Vis. Sci.
Identifiant (ID) PubMed	31185090 [20]

Liens

- [1] <http://okina.univ-angers.fr/j.kouassinzou/publications>
- [2] <http://okina.univ-angers.fr/j.chao/publications>
- [3] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=37852>
- [4] <http://okina.univ-angers.fr/s.leruez/publications>
- [5] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=37853>
- [6] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=1065>
- [7] <http://okina.univ-angers.fr/cbocca/publications>
- [8] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=30618>
- [9] <http://okina.univ-angers.fr/patrizia.bonneau/publications>
- [10] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=30621>
- [11] <http://okina.univ-angers.fr/d.bonneau/publications>
- [12] <http://okina.univ-angers.fr/gi.simard/publications>
- [13] <http://okina.univ-angers.fr/d.milea/publications>
- [14] <http://okina.univ-angers.fr/guy.lenaers/publications>
- [15] <http://okina.univ-angers.fr/v.procaccio/publications>
- [16] <http://okina.univ-angers.fr/pascal.reynier/publications>
- [17] <http://okina.univ-angers.fr/publications/ua19841>
- [18] <http://dx.doi.org/10.1167/iovs.19-27099>
- [19] <https://iovs.arvojournals.org/article.aspx?articleid=2736058>
- [20] <http://www.ncbi.nlm.nih.gov/pubmed/31185090?dopt=Abstract>

Publié sur *Okina* (<http://okina.univ-angers.fr>)